

3

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
24 October 2002 (24.10.2002)

PCT


(10) International Publication Number  
WO 02/083676 A1

- (51) International Patent Classification<sup>7</sup>: C07D 487/04, A01N 43/90 (74) Common Representative: BASF AKTIENGESELLSCHAFT; 67056 Ludwigshafen (DE).
- (21) International Application Number: PCT/EP02/03829 (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (22) International Filing Date: 6 April 2002 (06.04.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 01109011.5 11 April 2001 (11.04.2001) EP (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- (71) Applicant (*for all designated States except US*): BASF AKTIENGESELLSCHAFT [DE/DE]; 67056 Ludwigshafen (DE).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): TORMO I BLASCO, Jordi [ES/DE]; Mühlweg 47, 67117 Limburgerhof (DE). AMMERMAN, Eberhard [DE/DE]; Von-Gager-Str. 2, 64646 Heppenheim (DE). PEES, Klaus-Jürgen [DE/DE]; Soonwaldstr. 9, 55129 Mainz (DE). ALBERT, Guido [DE/DE]; Volxheimerstr. 4, 55546 Hackenheim (DE). REHNIG, Annerose [DE/DE]; Rathenaustasse 11, 55218 Ingelheim (DE). SEARCH, Debra [US/US]; 55 High Bridge Road, New Egypt, NJ 08533 (US).
- Published:**
- with international search report
  - before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 02/083676 A1

(54) Title: 5-HALOGEN-6-PHENYL-7-FLUORALKYLAMINO-TRIAZOLOPYRIMIDINES AS FUNGICIDES




(I)

(57) Abstract: 5-Halogen-6-phenyl-7-fluoroalkylamino-triazolopyrimidines of formula I: (I) in which R<sup>1</sup> is hydrogen, fluoro, alkyl, alkenyl, alkynyl, alkadienyl, where the carbon chains of these radicals may be unsubstituted or substituted as defined in the description; R<sup>2</sup> is hydrogen, alkyl, alkenyl, alkynyl, alkadienyl, where these radicals may be unsubstituted or substituted; R<sup>3</sup> is fluoroalkyl or fluoroalkenyl; X is halogen; n is 0 or an integer from 1 to 4; L each independently is halogen, nitro, alkyl, haloalkyl, alkoxy or haloalkoxy. Processes for their preparation, compositions containing them and to their use for combating phytopathogenic fungi.

## 5-Halogen-6-phenyl-7-fluoroalkylamino-triazolopyrimidines

## Description

The present Invention relates to 5-halogen-6-phenyl-7-fluoroalkylamino-triazolopyrimidines of formula I



I

in which

$R^1$  is hydrogen, fluoro,  $C_1$ - $C_{10}$ -alkyl,  $C_2$ - $C_{10}$ -alkenyl,  $C_2$ - $C_{10}$ -alkynyl,  $C_2$ - $C_{10}$ -alkadienyl,

where the carbon chains of these radicals may be unsubstituted or partially or fully halogenated or may carry one to three groups  $R^a$ ,

$R^a$  is cyano, nitro, hydroxyl,  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -haloalkyl,  $C_1$ - $C_6$ -alkylcarbonyl,  $C_3$ - $C_6$ -cycloalkyl,  $C_1$ - $C_6$ -alkoxy,  $C_1$ - $C_6$ -haloalkoxy,  $C_1$ - $C_6$ -alkoxycarbonyl,  $C_1$ - $C_6$ -alkylthio,  $C_1$ - $C_6$ -alkylamino,

di- $C_1$ - $C_6$ -alkylamino,  $C_2$ - $C_6$ -alkenyl,  $C_2$ - $C_6$ -alkenyloxy,  $C_3$ - $C_6$ -alkynyloxy and  $C_1$ - $C_4$ -alkylenedioxy, which may be halogenated;

$R^2$  is hydrogen,  $C_1$ - $C_{10}$ -alkyl,  $C_2$ - $C_{10}$ -alkenyl,  $C_2$ - $C_{10}$ -alkynyl,  $C_2$ - $C_{10}$ -alkadienyl, where these radicals may be unsubstituted or partially or fully halogenated or may carry one to three groups  $R^a$ ,

$R^3$  is  $C_2$ - $C_8$ -fluoroalkyl or  $C_2$ - $C_8$ -fluoroalkenyl;

X is halogen;

n is 0 or an integer from 1 to 4;

L each independently is halogen, nitro,  $C_1$ - $C_{10}$ -alkyl,  $C_1$ - $C_6$ -haloalkyl,  $C_1$ - $C_{10}$ -alkoxy or  $C_1$ - $C_6$ -haloalkoxy.

Moreover, the invention relates to processes for their preparation, compositions containing them and to their use for combating phytopathogenic fungi.

- 5 EP-A 071 792 discloses 6-phenyl-7-amino-triazolopyrimidines where the 5-position is substituted by hydrogen or alkyl or aryl groups.

- EP-A 550 113 relates to 5-H- and 5-halogen-6-phenyl-7-amino-tria-  
10 zolopyrimidines where the 7-amino group is further substituted.

WO-A 98/46607 discloses fungicidal triazolopyrimidines, which are substituted in the 6-position by a 2,4,6-trifluorophenyl group.

- 15 WO-A 98/46608 discloses fungicidal triazolopyrimidines, which are substituted in the 7-position by a fluoroalkylamino group.

The compounds disclosed in the documents discussed above are said to be active against various phytopathogenic fungi.

20

It is an object of the present invention to provide compounds having improved fungicidal activity.

- We have found that this object is achieved by the compounds de-  
25 fined at the outset. Furthermore, we have found processes for their preparation, compositions comprising them and methods for controlling phytopathogenic fungi using the compounds I.


- The compounds of the formula I differ from the compounds known  
30 from closest prior art, which is WO-A 98/46608, in the specific definition of the 7-fluoroalkylamino group, wherein the carbon chain in  $R^3$  consists of at least two carbon atoms.

- A 4- to 6- membered heterocyclic group may be any heterocyclic  
35 group with 4 to 6 ring atoms, interrupted by one or more heteroatoms selected from sulfur, nitrogen, and oxygen, preferably oxygen. A halogen atom suitable denotes a fluorine, chlorine or bromine atom.

- 40 The present invention further provides a process for the preparation of compounds of formula I as defined above which comprises treating a 5,7-dihalo compound of formula II in which X is halogen with an amine of formula III

45

3



The reaction between the 5,7-dihalo compound II and the amine of formula III can be carried out under conditions known from WO-A 98/46608.

10

The reaction is preferably carried out in the presence of a solvent. Suitable solvents include ethers, such as dioxane, diethyl ether and, especially, tetrahydrofuran, halogenated hydrocarbons such as dichloromethane and aromatic hydrocarbons, for example

15 toluene.

The reaction is suitably carried out at a temperature in the range from 0°C to 70°C, the preferred reaction temperature being from 10°C to 35°C.

20

It is also preferred that the reaction is carried out in the presence of a base. Suitable bases include tertiary amines, such as triethylamine, and inorganic bases, such as potassium carbonate or sodium carbonate. Alternatively, an excess of the compound

25 of formula III may serve as a base.

In a further embodiment of the invention the amine III is activated by complexation with a free or complexed metal atom, such as lithium, sodium or copper, and the reaction of II is carried

30

out with the complexed amine III'. In formula III' M denotes a free or complexed metal atom. The reaction conditions are in general similar as mentioned above.



40 The reaction mixtures are worked up in a customary manner, for example by mixing with water, phase separation and, if required, chromatographic purification of the crude products. Some of the end products are obtained in the form of colorless or slightly brownish, viscous oils, which are purified or freed from volatile

45

components under reduced pressure and at moderately elevated tem-

peratures. If the end products are obtained as solids, purification can also be carried out by recrystallization or digestion.

Compounds of formula II are known in the art and can be obtained  
5 by synthesis routes disclosed in EP-A 550 113 and WO-A 98/46608.

If individual compounds I are not obtainable by the routes described above, they can be prepared by derivatization of other compounds I.

10

In the symbol definitions given in the formulae above, collective terms were used which generally represent the following substituents:

15 - halogen: fluorine, chlorine, bromine and iodine;

- C<sub>1</sub>-C<sub>10</sub>-alkyl and the alkyl moieties of C<sub>1</sub>-C<sub>10</sub>-haloalkyl: saturated, straight-chain or branched hydrocarbon radicals having 1 to 10, especially 1 to 6 carbon atoms, for example C<sub>1</sub>-C<sub>4</sub>-alkyl as  
20 mentioned above or pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, 2,2-di-methylpropyl, 1-ethylpropyl, hexyl, 1,1-dimethylpropyl, 1,2-dimethylpropyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 1,3-dimethylbutyl, 2,2-dimethylbutyl, 2,3-dimethylbutyl,  
25 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethylbutyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl and 1-ethyl-2-methylpropyl;

- C<sub>1</sub>-C<sub>6</sub>-haloalkyl and the haloalkyl moieties of C<sub>1</sub>-C<sub>6</sub>-haloalkoxy:  
30 straight-chain or branched alkyl groups having 1 to 6, preferably 1 to 4 carbon atoms (as mentioned above), where the hydrogen atoms in these groups may be partially or fully replaced by halogen atoms as mentioned above, for example C<sub>1</sub>-C<sub>2</sub>-haloalkyl, such as chloromethyl, bromomethyl, dichloromethyl, trichloromethyl, fluoromethyl, difluoromethyl, trifluoromethyl, chlorofluoromethyl,  
35 dichlorofluoromethyl, chlorodifluoromethyl, 1-chloroethyl, 1-bromoethyl, 1-fluoroethyl, 2-fluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl, 2-chloro-2-fluoroethyl, 2-chloro-2,2-difluoroethyl, 2,2-dichloro-2-fluoroethyl, 2,2,2-trichloroethyl and  
40 pentafluoroethyl;

- C<sub>1</sub>-C<sub>6</sub>-fluoroalkyl and the fluoroalkyl moieties of C<sub>1</sub>-C<sub>6</sub>-fluoroalkoxy: straight-chain or branched alkyl groups having 1 to 6, preferably 1 to 4 carbon atoms (as mentioned above), where the  
45 hydrogen atoms in these groups may be partially or fully replaced by fluorine atoms as mentioned above, for example C<sub>1</sub>-C<sub>2</sub>-fluoroalkyl, such as fluoromethyl, difluoromethyl, trifluoromethyl,

1-fluoroethyl, 2-fluoroethyl, 2,2-difluoroethyl, 2,2,2-trifluoroethyl and pentafluoroethyl;

- C<sub>2</sub>-C<sub>10</sub>-alkenyl: unsaturated, straight-chain or branched hydro-  
5 carbon radicals having 2 to 10, especially 3 to 6 carbon atoms and a double bond in any position, for example 1-propenyl, 2-propenyl, 1-methylethenyl, 1-butenyl, 2-butenyl, 3-butenyl, 1-methyl-1-propenyl, 2-methyl-1-propenyl, 1-methyl-2-propenyl and 2-methyl-2-propenyl;

10

C<sub>2</sub>-C<sub>10</sub>-alkynyl: straight-chain or branched hydrocarbon radicals having 2 to 10, especially 2 to 4 carbon atoms and a triple bond in any position, for example ethynyl, 1-propynyl, 2-propynyl, 1-butyne, 2-butyne, 3-butyne and 1-methyl-2-propynyl.

15

With respect to their intended use, preference is given to triazolopyrimidines of the formula I having the following substituents, where the preference is valid in each case on its own or in combination:

20

The particularly preferred embodiments of the intermediates with respect to the variables correspond to those of the radicals X, R<sup>1</sup> and R<sup>2</sup> of formula I.

25 A preferred alkyl moiety is an ethyl or especially a methyl group.

A preferred haloalkyl moiety is the 2,2,2-trifluoroethyl or 1,1,1-trifluoroprop-2-yl group;

30

A preferred alkenyl moiety is allyl or especially a 2-methylallyl group.

Likewise, preference is given to compounds of formula I wherein R<sup>1</sup>  
35 is not hydrogen.

Compounds of formula I are preferred in which R<sup>1</sup> represents fluorine or straight-chained or branched C<sub>1</sub>-C<sub>10</sub>-alkyl, in particular branched C<sub>3</sub>-C<sub>10</sub>-alkyl.

40

Furthermore, particular preference is given to compounds I in which R<sup>1</sup> is fluorine.

Particular preference is given to compounds I in which R<sup>2</sup> represents a hydrogen atom, a C<sub>1</sub>-C<sub>10</sub>-alkyl or a C<sub>1</sub>-C<sub>10</sub>-haloalkyl group,  
45 in particular a hydrogen atom.

Besides, particular preference is given to compounds I in which R<sup>2</sup> is hydrogen.

Moreover, particular preference is given to compounds I in which  
5 R<sup>2</sup> is methyl.

Particular preference is given to compounds of formula I in which R<sup>3</sup> denotes a group A

10



wherein

15 Y<sup>1</sup> is hydrogen, fluoro or C<sub>1</sub>-C<sub>6</sub>-fluoroalkyl,

Y<sup>2</sup> is hydrogen or fluoro, or

Y<sup>1</sup> and Y<sup>2</sup> taken together form a double bond; and

20


m is 0 or 1.

If R<sup>3</sup> denotes a C<sub>1</sub>-C<sub>10</sub>-haloalkyl group, preferably a polyfluorinated alkyl group, in particular a 2,2,2-trifluoroethyl, a


25 2-(1,1,1-trifluoropropyl) or a 2-(1,1,1-trifluorobutyl) group, R<sup>2</sup> preferably represents a hydrogen atom.

L<sub>n</sub> preferably is halogen or C<sub>1</sub>-C<sub>6</sub>-alkoxy. A preferred embodiment are compounds of formula I in which

30



represents



35


wherein # denotes the link to the triazolopyrimidine moiety, L<sup>1</sup> through L<sup>4</sup> each independently represent hydrogen, especially fluo-  
rine, chlorine, methyl or methoxy, in particular wherein L<sup>1</sup> is  
fluoro, L<sup>2</sup> is hydrogen or fluoro, L<sup>3</sup> is hydrogen or fluoro or me-

40 thoxy and L<sup>4</sup> denotes hydrogen, fluoro chloro or methyl.

Moreover, particular preference is given to compounds of the for-  
mula I in which n is 2 or 3. Most preferred L<sup>4</sup> is not hydrogen.

45 Furthermore, particular preference is given to compounds of for-  
mula IA in which R<sup>1</sup> to R<sup>3</sup> have the meaning as defined in formula  
I and L<sup>1</sup> to L<sup>4</sup> have the meanings given above.

7



IA

Particular preference is given to compounds of formula IA in which L<sup>1</sup> denotes halogen and L<sup>3</sup> and L<sup>4</sup> each independently represent hydrogen, halogen or C<sub>1</sub>-C<sub>4</sub>-alkoxy.

Most preferred are compounds of formula IA in which L<sup>1</sup> is fluoro, L<sup>3</sup> is hydrogen and L<sup>4</sup> is chloro.

Particularly preference is given to compounds of formula IA wherein R<sup>2</sup> is hydrogen, L<sup>1</sup> and L<sup>4</sup> independently represent fluorine or chlorine atoms, and L<sup>3</sup> denotes hydrogen, fluorine, chlorine or methoxy.

Included in the scope of the present invention are (R) and (S) isomers of compounds of general formula I having a chiral center and the racemates thereof, and salts, N-oxides and acid addition compounds.

With respect to their use, particular preference is given to the compounds I compiled in the tables below. The groups mentioned in the table for a substituent are furthermore for their part, independently of the combination in which they are mentioned, a particularly preferred embodiment of the respective substituents.

30

Table 1

Compounds of the formula IA, in which L<sup>1</sup> is fluoro, L<sup>3</sup> is hydrogen, L<sup>4</sup> is chloro and R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> correspond to one row in Table A

35

Table 2

Compounds of the formula IA, in which L<sup>1</sup> and L<sup>4</sup> each are fluoro, L<sup>3</sup> is hydrogen and R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> correspond to one row in Table A

40 Table 3

Compounds of the formula IA, in which L<sup>1</sup> and L<sup>4</sup> each are chloro, L<sup>3</sup> is hydrogen and R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> correspond to one row in Table A

45



Table 4

Compounds of the formula IA, in which L<sup>1</sup> is methyl, L<sup>3</sup> is hydrogen, L<sup>4</sup> is fluoro and R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> correspond to one row in Table A

5

Table 5

Compounds of the formula IA, in which L<sup>1</sup>, L<sup>3</sup> and L<sup>4</sup> each are fluoro and R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> correspond to one row in Table A

10 Table 6

Compounds of the formula IA, in which L<sup>1</sup> and L<sup>4</sup> each are fluoro, L<sup>3</sup> is methoxy and R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> correspond to one row in Table A

Table 7

15 Compounds of the formula IA, in which L<sup>1</sup> and L<sup>3</sup> are each hydrogen, L<sup>4</sup> is fluoro and R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> correspond to one row in Table A

Table 8

Compounds of the formula IA, in which L<sup>1</sup> and L<sup>3</sup> are each hydrogen,  
20 L<sup>4</sup> is chloro and R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> correspond to one row in Table A

Table 9


Compounds of the formula I, in which L<sub>n</sub> is 2-F-5-NO<sub>2</sub>, X is chloro and R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> correspond to one row in Table A

25

Table 10

Compounds of the formula I, in which L<sub>n</sub> is pentafluoro, X is chloro and R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> correspond to one row in Table A

30 Table A



IA

35

No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>
A-1	F	H	CH <sub>2</sub> CF <sub>3</sub>
40 A-2	F	CH <sub>3</sub>	CH <sub>2</sub> CF <sub>3</sub>
A-3	F	CH <sub>2</sub> CH <sub>3</sub>	CH <sub>2</sub> CF <sub>3</sub>
A-4	F	CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	CH <sub>2</sub> CF <sub>3</sub>
A-5	F	H	CH <sub>2</sub> C(CF <sub>3</sub> )=CH <sub>2</sub>
45 A-6	F	CH <sub>3</sub>	CH <sub>2</sub> C(CF <sub>3</sub> )=CH <sub>2</sub>
A-7	F	CH <sub>2</sub> CH <sub>3</sub>	CH <sub>2</sub> C(CF <sub>3</sub> )=CH <sub>2</sub>
A-8	F	H	CH <sub>2</sub> C(CH <sub>3</sub> )=CF <sub>2</sub>

No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>
A-9	F	CH <sub>3</sub>	CH <sub>2</sub> C (CH <sub>3</sub> ) =CF <sub>2</sub>
A-10	F	CH <sub>2</sub> CH <sub>3</sub>	CH <sub>2</sub> C (CH <sub>3</sub> ) =CF <sub>2</sub>
5 A-11	F	H	CH <sub>2</sub> CF=CH <sub>2</sub>
A-12	F	CH <sub>3</sub>	CH <sub>2</sub> CF=CH <sub>2</sub>
A-13	F	CH <sub>2</sub> CH <sub>3</sub>	CH <sub>2</sub> CF=CH <sub>2</sub>
A-14	F	H	CH <sub>2</sub> CH=CF <sub>2</sub>
10 A-15	F	CH <sub>3</sub>	CH <sub>2</sub> CH=CF <sub>2</sub>
A-16	F	CH <sub>2</sub> CH <sub>3</sub>	CH <sub>2</sub> CH=CF <sub>2</sub>
A-17	H	H	CF <sub>2</sub> CF <sub>3</sub>
A-18	H	CH <sub>3</sub>	CF <sub>2</sub> CF <sub>3</sub>
A-19	H	CH <sub>2</sub> CH <sub>3</sub>	CF <sub>2</sub> CF <sub>3</sub>
15 A-20	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	CF <sub>2</sub> CF <sub>3</sub>
A-21	H	H	CHFCF <sub>3</sub>
A-22	H	CH <sub>3</sub>	CHFCF <sub>3</sub>
A-23	H	CH <sub>2</sub> CH <sub>3</sub>	CHFCF <sub>3</sub>
20 A-24	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	CHFCF <sub>3</sub>
A-25	H	H	CH <sub>2</sub> CF <sub>2</sub> CF <sub>3</sub>
A-26	H	CH <sub>3</sub>	CH <sub>2</sub> CF <sub>2</sub> CF <sub>3</sub>
A-27	H	CH <sub>2</sub> CH <sub>3</sub>	CH <sub>2</sub> CF <sub>2</sub> CF <sub>3</sub>
25 A-28	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	CH <sub>2</sub> CF <sub>2</sub> CF <sub>3</sub>
A-29	H	H	CH (CH <sub>3</sub> ) CH <sub>2</sub> CF <sub>3</sub>
A-30	H	CH <sub>3</sub>	CH (CH <sub>3</sub> ) CH <sub>2</sub> CF <sub>3</sub>
A-31	H	CH <sub>2</sub> CH <sub>3</sub>	CH (CH <sub>3</sub> ) CH <sub>2</sub> CF <sub>3</sub>
30 A-32	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	CH (CH <sub>3</sub> ) CH <sub>2</sub> CF <sub>3</sub>
A-33	H	H	CH <sub>2</sub> CH=CF <sub>2</sub>
A-34	H	CH <sub>3</sub>	CH <sub>2</sub> CH=CF <sub>2</sub>
A-35	H	CH <sub>2</sub> CH <sub>3</sub>	CH <sub>2</sub> CH=CF <sub>2</sub>
A-36	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	CH <sub>2</sub> CH=CF <sub>2</sub>
35 A-37	H	H	CH <sub>2</sub> CF=CF <sub>2</sub>
A-38	H	CH <sub>3</sub>	CH <sub>2</sub> CF=CF <sub>2</sub>
A-39	H	CH <sub>2</sub> CH <sub>3</sub>	CH <sub>2</sub> CF=CF <sub>2</sub>
A-40	H	CH <sub>2</sub> CH <sub>2</sub> CH <sub>3</sub>	CH <sub>2</sub> CF=CF <sub>2</sub>

40 The compounds I are suitable as fungicides. They have outstanding activity against a broad spectrum of phytopathogenic fungi, in particular from the classes of the *Ascomycetes*, *Deuteromycetes*, *Phycomycetes* and *Basidiomycetes*. Some of them act systemically, and they can be employed in crop protection as foliar- and soil-  
 45 acting fungicides.

## 10

They are especially important for controlling a large number of fungi on a variety of crop plants such as wheat, rye, barley, oats, rice, maize, grass, bananas, cotton, soya, coffee, sugar cane, grapevines, fruit species, ornamentals and vegetables such as cucumbers, beans, tomatoes, potatoes and cucurbits, and on the seeds of these plants.

Specifically, they are suitable for controlling the following plant diseases:

- 10 Alternaria species on vegetables and fruit,  
Botrytis cinerea (gray mold) on strawberries, vegetables, ornamentals and grapevines,  
Cercospora arachidicola on peanuts,  
Erysiphe cichoracearum and Sphaerotheca fuliginea on cucurbits,
- 15 Blumeria graminis (powdery mildew) on cereals,  
Fusarium and Verticillium species on various plants,  
Helminthosporium species on cereals,  
Mycosphaerella species on bananas and peanuts,  
Phytophthora infestans on potatoes and tomatoes,
- 20 Plasmopara viticola on grapevines,  
Podosphaera leucotricha on apples,  
Pseudocercospora herpotrichoides on wheat and barley,  
Pseudoperonospora species on hops and cucumbers,  
Puccinia species on cereals,
- 25 Pyricularia oryzae on rice,  
Rhizoctonia species on cotton, rice and lawns,  
Septoria species in cereals,  
Uncinula necator on grapevines,  
Ustilago species on cereals and sugar cane, and
- 30 Venturia species (scab) on apples and pears.

Moreover, the compounds I are suitable for controlling harmful fungi such as Paecilomyces variotii in the protection of materials (e.g. wood, paper, paint dispersions, fibers and tissues) and in the protection of stored products.

The compounds I are applied by treating the fungi, or the plants, seeds, materials or the soil to be protected against fungal infection, with a fungicidally active amount of the active ingredients. Application can be effected both before and after infection of the materials, plants or seeds by the fungi.

In general, the fungicidal compositions comprise from 0.1 to 95, preferably 0.5 to 90, % by weight of active ingredient.

When used in crop protection, the rates of application are from 0.01 to 2.0 kg of active ingredient per ha, depending on the nature of the effect desired.

- 5 In the treatment of seed, amounts of active ingredient of from 0.001 to 0.1 g, preferably 0.01 to 0.05 g, are generally required per kilogram of seed.

- When used in the protection of materials or stored products, the  
10 rate of application of active ingredient depends on the nature of the field of application and on the effect desired. Rates of application conventionally used in the protection of materials are, for example, from 0.001 g to 2 kg, preferably 0.005 g to 1 kg, of active ingredient per cubic meter of material treated.

- 15 The compounds I can be converted into the customary formulations, e.g. solutions, emulsions, suspensions, dusts, powders, pastes and granules. The use form depends on the particular purpose; in any case, it should guarantee a fine and uniform distribution of  
20 the compound according to the invention.

- The formulations are prepared in a known manner, e.g. by extending the active ingredient with solvents and/or carriers, if desired using emulsifiers and dispersants, it also being possible  
25 to use other organic solvents as auxiliary solvents if water is used as the diluent. Auxiliaries which are suitable are essentially: solvents such as aromatics (e.g. xylene), chlorinated aromatics (e.g. chlorobenzenes), paraffins (e.g. mineral oil fractions), alcohols (e.g. methanol, butanol), ketones (e.g. cy-  
30 clohexanone), amines (e.g. ethanolamine, dimethylformamide) and water; carriers such as ground natural minerals (e.g. kaolins, clays, talc, chalk) and ground synthetic minerals (e.g. highly-disperse silica, silicates); emulsifiers such as non-ionic and anionic emulsifiers (e.g. polyoxyethylene fatty alcohol ethers,  
35 alkylsulfonates and arylsulfonates) and dispersants such as lignin-sulfite waste liquors and methylcellulose.

- Suitable surfactants are alkali metal, alkaline earth metal and ammonium salts of lignosulfonic acid, naphthalenesulfonic acid,  
40 phenolsulfonic acid, dibutyl-naphthalenesulfonic acid, alkylaryl-sulfonates, alkyl sulfates, alkylsulfonates, fatty alcohol sulfates and fatty acids and their alkali metal and alkaline earth metal salts, salts of sulfated fatty alcohol glycol ether, condensates of sulfonated naphthalene and naphthalene derivatives  
45 with formaldehyde, condensates of naphthalene or of naphthalene-sulfonic acid with phenol or formaldehyde, polyoxyethylene octyl-phenyl ether, ethoxylated isooctylphenol, octylphenol, nonylphe-

nol, alkylphenol polyglycol ethers, tributylphenyl polyglycol ethers, alkylaryl polyether alcohols, isotridecyl alcohol, fatty alcohol/ethylene oxide condensates, ethoxylated castor oil, polyoxyethylene alkyl ethers, ethoxylated polyoxypropylene, lauryl  
5 alcohol polyglycol ether acetal, sorbitol esters, lignin-sulfite waste liquors and methylcellulose.

- Substances which are suitable for the preparation of directly sprayable solutions, emulsions, pastes or oil dispersions are
- 10 mineral oil fractions of medium to high boiling point, such as kerosene or diesel oil, furthermore coal tar oils and oils of vegetable or animal origin, aliphatic, cyclic and aromatic hydrocarbons, e.g. benzene, toluene, xylene, paraffin, tetrahydronaphthalene, alkylated naphthalenes or their derivatives, methanol,  
15 ethanol, propanol, butanol, chloroform, carbon tetrachloride, cyclohexanol, cyclohexanone, chlorobenzene, isophorone, strongly polar solvents, e.g. dimethylformamide, dimethyl sulfoxide, N-methylpyrrolidone and water.
- 20 Powders, materials for scattering and dusts can be prepared by mixing or concomitantly grinding the active substances with a solid carrier.

- Granules, e.g. coated granules, impregnated granules and homogeneous granules, can be prepared by binding the active ingredients to solid carriers. Examples of solid carriers are mineral earths, such as silicas, silica gels, silicates, talc, kaolin, attaclay, limestone, lime, chalk, bole, loess, clay, dolomite, diatomaceous earth, calcium sulfate, magnesium sulfate, magnesium oxide,  
25 ground synthetic materials, fertilizers, e.g. ammonium sulfate, ammonium phosphate, ammonium nitrate, ureas, and products of vegetable origin, such as cereal meal, tree bark meal, wood meal and nutshell meal, cellulose powders and other solid carriers.
- 30 In general, the formulations comprise of from 0.01 to 95% by weight, preferably from 0.1 to 90% by weight, of the active ingredient. The active ingredients are employed in a purity of from 90% to 100%, preferably 95% to 100% (according to NMR spectrum).

- 40 The following are exemplary formulations:

- I. 5 parts by weight of a compound according to the invention are mixed intimately with 95 parts by weight of finely divided kaolin. This gives a dust which comprises 5% by weight  
45 of the active ingredient.

- II. 30 parts by weight of a compound according to the invention are mixed intimately with a mixture of 92 parts by weight of pulverulent silica gel and 8 parts by weight of paraffin oil which had been sprayed onto the surface of this silica gel. This gives a formulation of the active ingredient with good adhesion properties (comprises 23% by weight of active ingredient).
- III. 10 parts by weight of a compound according to the invention are dissolved in a mixture composed of 90 parts by weight of xylene, 6 parts by weight of the adduct of 8 to 10 mol of ethylene oxide and 1 mol of oleic acid N-monoethanolamide, 2 parts by weight of calcium dodecylbenzenesulfonate and 2 parts by weight of the adduct of 40 mol of ethylene oxide and 1 mol of castor oil (comprises 9% by weight of active ingredient).
- IV. 20 parts by weight of a compound according to the invention are dissolved in a mixture composed of 60 parts by weight of cyclohexanone, 30 parts by weight of isobutanol, 5 parts by weight of the adduct of 7 mol of ethylene oxide and 1 mol of isooctylphenol and 5 parts by weight of the adduct of 40 mol of ethylene oxide and 1 mol of castor oil (comprises 16% by weight of active ingredient).
- V. 80 parts by weight of a compound according to the invention are mixed thoroughly with 3 parts by weight of sodium diisobutylphenylthalethene- $\alpha$ -sulfonate, 10 parts by weight of the sodium salt of a lignosulfonic acid from a sulfite waste liquor and 7 parts by weight of pulverulent silica gel, and the mixture is ground in a hammer mill (comprises 80% by weight of active ingredient).
- VI. 90 parts by weight of a compound according to the invention are mixed with 10 parts by weight of N-methyl- $\alpha$ -pyrrolidone, which gives a solution which is suitable for use in the form of microdrops (comprises 90% by weight of active ingredient).
- VII. 20 parts by weight of a compound according to the invention are dissolved in a mixture composed of 40 parts by weight of cyclohexanone, 30 parts by weight of isobutanol, 20 parts by weight of the adduct of 7 mol of ethylene oxide and 1 mol of isooctylphenol and 10 parts by weight of the adduct of 40 mol of ethylene oxide and 1 mol of castor oil. Pouring the solution into 100,000 parts by weight of water and finely

distributing it therein gives an aqueous dispersion which comprises 0.02% by weight of the active ingredient.

- VIII. 20 parts by weight of a compound according to the invention  
5 are mixed thoroughly with 3 parts by weight of sodium diisobutyl-naphthalene- $\alpha$ -sulfonate, 17 parts by weight of the sodium salt of a lignosulfonic acid from a sulfite waste liquor and 60 parts by weight of pulverulent silica gel, and the mixture is ground in a hammer mill. Finely distributing  
10 the mixture in 20,000 parts by weight of water gives a spray mixture which comprises 0.1% by weight of the active ingredient.

The active ingredients can be used as such, in the form of their  
15 formulations or the use forms prepared therefrom, e.g. in the form of directly sprayable solutions, powders, suspensions or dispersions, emulsions, oil dispersions, pastes, dusts, materials for spreading, or granules, by means of spraying, atomizing, dusting, scattering or pouring. The use forms depend entirely on  
20 the intended purposes; in any case, this is intended to guarantee the finest possible distribution of the active ingredients according to the invention.

Aqueous use forms can be prepared from emulsion concentrates,  
25 pastes or wettable powders (sprayable powders, oil dispersions) by adding water. To prepare emulsions, pastes or oil dispersions, the substances as such or dissolved in an oil or solvent, can be homogenized in water by means of wetter, tackifier, dispersant or emulsifier. Alternatively, it is possible to prepare concentrates  
30 composed of active substance, wetter, tackifier, dispersant or emulsifier and, if appropriate, solvent or oil, and such concentrates are suitable for dilution with water.

The active ingredient concentrations in the ready-to-use products  
35 can be varied within substantial ranges. In general, they are from 0.0001 to 10%, preferably from 0.01 to 1%.

The active ingredients may also be used successfully in the ultra-low-volume process (ULV), it being possible to apply formula-  
40 tions comprising over 95% by weight of active ingredient, or even the active ingredient without additives.

Various types of oils, herbicides, fungicides, other pesticides, or bactericides may be added to the active ingredients, if ap-  
45 propriate also only immediately prior to use (tank mix). These agents can be admixed with the agents according to the invention in a weight ratio of 1:10 to 10:1.

In the use form as fungicides, the compositions according to the invention can also be present together with other active ingredients, e.g. with herbicides, insecticides, growth regulators, fungicides or else with fertilizers. Mixing the compounds I or the  
 5 compositions comprising them in the use form as fungicides with other fungicides frequently results in a broader fungicidal spectrum of action.

The following list of fungicides, together with which the com-  
 10 pounds according to the invention can be used, is intended to illustrate the possible combinations, but not to impose any limitation:

15 sulfur, dithiocarbamates and their derivatives, such as iron(III) dimethyldithiocarbamate, zinc dimethyldithiocarbamate, zinc ethylenebisdithiocarbamate, manganese ethylenebisdithiocarbamate, manganese zinc ethylenediaminebisdithiocarbamate, tetramethylthiuram disulfide, ammonia complex of zinc  
 20 (N,N-ethylenebisdithiocarbamate), ammonia complex of zinc (N,N'-propylenebisdithiocarbamate), zinc (N,N'-propylenebisdithiocarbamate), N,N'-polypropylenebis(thiocarbamoyl)disulfide;  
 nitro derivatives, such as dinitro(1-methylheptyl)phenyl crotonate, 2-sec-butyl-4,6-dinitrophenyl 3,3-dimethylacrylate,  
 25 2-sec-butyl-4,6-dinitrophenylisopropyl carbonate, diisopropyl 5-nitro-isophthalate;  
 heterocyclic substances, such as 2-heptadecyl-2-imidazoline acetate, 2,4-dichloro-6-(o-chloroanilino)-s-triazine, O,O-diethyl phthalimidophosphonothioate, 5-amino-1-[bis(dimethylamino)phosphinyl]-3-phenyl-1,2,4- triazole, 2,3-dicyano-1,4-dithioanthraquinone, 2-thio-1,3-dithiolo[4,5-b]quinoxaline, methyl 1-(butylcarbamoyle)-2-benzimidazolecarbamate,  
 30 2-methoxycarbonylaminobenzimidazole, 2-(2-furyl)benzimidazole, 2-(4-thiazolyl)benzimidazole, N-(1,1,2,2-tetrachloroethylthio)tetrahydrophthalimide, N-trichloromethylthiotetrahydrophthalimide, N-trichloromethylthiophthalimide,  
 35 N-dichlorofluoromethylthio-N',N'-dimethyl-N-phenylsulfo- diamide, 5-ethoxy-3-trichloromethyl-1,2,3-thiadiazole, 2-thiocyanatomethylthiobenzothiazole, 1,4-dichloro-2,5-dimethoxybenzene, 4-(2-chlorophenylhydrazono)-3-methyl-5-isoxazolone,  
 40 pyridine-2-thiol 1-oxide, 8-hydroxyquinoline or its copper salt, 2,3-dihydro-5-carboxanilido-6-methyl-1,4-oxathiine, 2,3-dihydro-5-carboxanilido-6-methyl-1,4-oxathiine 4,4-dioxide, 2-methyl-5,6-dihydro-4H-pyran-3-carboxanilide, 2-methylfuran-3-carboxanilide, 2,5-dimethylfuran-3-carboxanilide,  
 45 2,4,5-trimethylfuran-3-carboxanilide, N-cyclohexyl- 2,5-dimethylfuran-3-carboxamide, N-cyclohexyl-N-methoxy-2,5-dimethyl-



furan-3-carboxamide, 2-methylbenzanilide, 2-iodobenzanilide, N-formyl-N-morpholine-2,2,2-trichloroethyl acetal, piperazine-1,4-diylbis-1-(2,2,2-trichloroethyl)formamide, 1-(3,4-dichloroanilino)-1-formylamino-2,2,2-trichloroethane;

5 2,6-dimethyl-N-tridecylmorpholine or its salts, 2,6-dimethyl-N-cyclododecylmorpholine or its salts, N-[3-(p-tert-butylphenyl)-2-methylpropyl]-cis-2,6-dimethyl-morpholine, N-[3-(p-tert-butylphenyl)-2-methylpropyl]-piperidine, 1-[2-(2,4-dichlorophenyl)-4-ethyl-1,3-dioxolan-2-yl-ethyl]-1H-1,2,4-triazole, 1-[2-(2,4-dichlorophenyl)-4-n-propyl-1,3-dioxolan-2-yl-ethyl]-1H-1,2,4-triazole, N-(n-propyl)-N-(2,4,6-trichlorophenoxyethyl)-N'-imidazolyl-urea, 1-(4-chlorophenoxy)-3,3-dimethyl-1-(1H-1,2,4-triazol-1-yl)-2-butanone, 1-(4-chlorophenoxy)-3,3-dimethyl-1-(1H-1,2,4-triazol-1-yl)-2-butanol, (2RS,3RS)-1-[3-(2-chlorophenyl)-2-(4-fluorophenyl)-oxiran-2-ylmethyl]-1H-1,2,4-triazole,  $\alpha$ -(2-chlorophenyl)- $\alpha$ -(4-chlorophenyl)-5-pyrimidinemethanol, 5-butyl-2-dimethylamino-4-hydroxy-6-methylpyrimidine, bis(p-chlorophenyl)-3-pyridinemethanol, 1,2-bis(3-ethoxycarbonyl-2-thioureido)benzene, 1,2-bis(3-methoxycarbonyl-2-thioureido)benzene, strobilurines such as azoxystrobin, kresoxim methyl, methyl-E-methoxyimino-[ $\alpha$ -(2-phenoxyphenyl)]-acetamide, methyl

25 E-methoxyimino-[ $\alpha$ -(2,5-dimethylphenoxy)-o-tolyl]acetamide, picoxystrobin, pyraclostrobin, trifloxystrobin, anilinopyrimidines such as N-(4,6-dimethylpyrimidin-2-yl)aniline, N-[4-methyl-6-(1-propynyl)pyrimidin-2-yl]-aniline, N-[4-methyl-6-cyclopropylpyrimidin-2-yl]aniline,

30 phenylpyrroles such as 4-(2,2-difluoro-1,3-benzodioxol-4-yl)pyrrole-3-carbonitrile, cinnamamides such as 3-(4-chlorophenyl)-3-(3,4-dimethoxyphenyl)acryloylmorpholine, and a variety of fungicides such as dodecylguanidine acetate,

35 3-[3-(3,5-dimethyl-2-oxycyclohexyl)-2-hydroxyethyl]glutaramide, hexachlorobenzene, methyl N-(2,6-dimethylphenyl)-N-(2-furoyl)-DL-alaninate, DL-N-(2,6-dimethylphenyl)-N-(2'-methoxyacetyl)-alanine methyl ester, N-(2,6-dimethylphenyl)-N-chloroacetyl-D,L-2-amino-butylolactone,

40 DL-N-(2,6-dimethylphenyl)-N-(phenylacetyl)alanine methyl ester, 5-methyl-5-vinyl-3-(3,5-dichlorophenyl)-2,4-dioxo-1,3-oxazolidine, 3-[3,5-dichlorophenyl(5-methyl-5-methoxymethyl)-1,3-oxazolidine-2,4-dione, 3-(3,5-dichlorophenyl)-1-isopropylcarbamoylethylhydantoin, N-(3,5-dichlorophenyl)-1,2-dimethylcyclopropane-1,2-dicarboximide, 2-cyano-[N-(ethylaminocarbonyl)-2-methoximino]acetamide, 1-[2-(2,4-dichlorophenyl)pentyl]-1H-1,2,4-triazole, 2,4-di-

45

fluoro- $\alpha$ -(1H-1,2,4-triazolyl-1-methyl)benzhydryl alcohol,  
 N-(3-chloro-2,6-dinitro-4-trifluoromethylphe-  
 nyl)-5-trifluoromethyl-3-chloro-2-aminopyridine,  
 1-((bis(4-fluorophenyl)methylsilyl)methyl)-1H-1,2,4-triazole.

5

### Synthesis Examples


With due modification of the starting compounds, the protocols shown in the synthesis examples below were used for obtaining  
 10 further compounds I. The resulting compounds, together with physical data, are listed in the Table which follows.

Example 1: Preparation of 5-chloro-6-(2-chloro-6-fluorophe-  
 nyl)-7-(1,1,1-trifluorobut-4-yl)-amino-[1,2,4]-triazolo[1,5- $\alpha$ ]py-  
 15 rimidine [I-9]

A solution of 1.5 mmol 1,1,1-trifluorobutane-4-amine, 1.5 mmol triethylamine in 10 ml dichloromethane is added to a solution of 1.5 mmol 5,7-dichloro-6-(2-chloro-6-fluorophenyl)-[1,2,4]-tri-  
 20 azolo[1,5- $\alpha$ ]pyrimidine [cf. EP-A 550 113] in 20 ml dichloromethane. The combined solution was stirred for about 16 hours at 20 to 25°C and subsequently washed with diluted (5%) hydrochloric acid. The organic phase was separated, dried and filtered. Distillative removal of the solvent from the filtrate and  
 25 chromatography over silica gel yielded 0.45 g of the title compound of mp. 115°C.

Table I

30



35

No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	L <sub>n</sub>	phys. data (m.p. [°C])
I-1	H	H	CHFCF <sub>3</sub>	2,4,6-F <sub>3</sub>	189
I-2	H	H	CH <sub>2</sub> CF=CF <sub>2</sub>	2,4,6-F <sub>3</sub>	137
40 I-3	H	H	C(CH <sub>3</sub> )CH <sub>2</sub> CF <sub>3</sub>	2,4,6-F <sub>3</sub>	143
I-4	F	H	CH <sub>2</sub> CF <sub>3</sub>	F <sub>5</sub>	200
I-5	H	H	CH <sub>2</sub> CF <sub>3</sub>	2,4,6-F <sub>3</sub>	202
I-6	H	H	CF <sub>2</sub> CF <sub>3</sub>	2-Cl-6-F	140
45 I-7	H	H	CH <sub>2</sub> CF <sub>3</sub>	2,6-F <sub>2</sub> -4-OCH <sub>3</sub>	161
I-8	H	H	CF <sub>2</sub> CF <sub>2</sub> CF <sub>3</sub>	2-Cl-6-F	182
I-9	H	H	CH <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	2-Cl-6-F	115

No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	L <sub>n</sub>	phys. data (m.p. [°C])
I-10	H	H	CH <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	2,4,6-F <sub>3</sub>	142
I-11	H	H	CH <sub>2</sub> CF=CF <sub>2</sub>	2,6-F <sub>2</sub>	130
I-12	H	H	CH <sub>2</sub> CF=CF <sub>2</sub>	2-Cl	129
I-13	H	H	CH <sub>2</sub> CF=CF <sub>2</sub>	2-F	111
I-14	H	H	CH <sub>2</sub> CF=CF <sub>2</sub>	2-Cl-6-F	107
I-15	H	H	CH <sub>2</sub> CF=CF <sub>2</sub>	F <sub>5</sub>	118
I-16	H	H	CHF <sub>2</sub> CF <sub>3</sub>	2-NO <sub>2</sub> -5-F	syrup
I-17	H	H	CHF <sub>2</sub> CF <sub>3</sub>	2-Cl-6-F	199
I-18	H	H	CHF <sub>2</sub> CF <sub>3</sub>	2,6-F <sub>2</sub>	80
I-19	H	H	CH <sub>2</sub> CH <sub>2</sub> CF <sub>3</sub>	2-F	81

15

Examples of the action against harmful fungi

20

The fungicidal action of the compounds of the formula I was demonstrated by the following experiments:

25

The active compounds, separately or together, were formulated as a 10% emulsion in a mixture of 70% by weight of cyclohexanone, 20% by weight of Nekanil® LN (Lutensol® AP6, wetting agent having emulsifying and dispersant action based on ethoxylated alkylphenols) and 10% by weight of Wettol® EM (nonionic emulsifier based on ethoxylated castor oil) and diluted with water to the desired concentration.

30

Use Example 1: Fungicidal control of apple scab  
(*Venturia inaequalis*)

35

Young apple seedlings of the cultivar "Common" were grown in pots to the 4 to 5 leaf stage. These plants were sprayed to run-off with an aqueous suspension, containing the concentration of active ingredient mentioned below, prepared from a stock solution containing 5 % of the active ingredient, 94 % cyclohexanone and 1 % emulsifier (Tween 20). After the plants had dried (3-5h), they were inoculated with an aqueous spore suspension of *Venturia inaequalis*. Then the trial plants were immediately transferred to a humid chamber with 22 to 24°C and a relative humidity close to 100 % and were cultivated there for 2 days. For a period of further 2 weeks a cultivation in a greenhouse followed at 21 to 23°C and a relative humidity about 95 %. Then the extent of fungal attack on the leaves was visually assessed as % diseased leaf area.

45

In this test, the plants which had been treated with 200 ppm of compounds I-1 to I-6 and I-8 to I-19 showed an infection of up to 15%, whereas the untreated plants were infected to 80%.

5 Use Example 2: Fungicidal control of early blight on tomatoes  
(*Alternaria solani*)

Young tomato seedlings of the cultivar "Pixie II" were grown in pots to the 2 to 4 leaf stage. These plants were sprayed to run-  
10 off with an aqueous suspension, containing the concentration of active ingredient mentioned below, prepared from a stock solution containing 5 % of the active ingredient, 94 % cyclohexanone and 1 % emulsifier (Tween 20). After the plants had dried (3-5h), they were inoculated with an aqueous spore suspension of *Alternaria*  
15 *solani* containing  $15 \times 10^3$  spores per ml. Then the trial plants were immediately transferred to a humid chamber with 22 to 24°C and a relative humidity close to 100 % for 36 h. For a period of further 2 to 3 days a cultivation in a greenhouse followed at 21 to 23°C and a relative humidity about 95 %. Then the extent of  
20 fungal attack on the leaves was visually assessed as % diseased leaf area.

In this test, the plants which had been treated with 200 ppm of compounds I-1 to I-7, I-9 to I-13, I-15, I-17, and I-18 showed an  
25 infection of up to 15%, whereas the untreated plants were infected to 90%.

Use Example 3: Fungicidal control of leaf spot on beets  
(*Cercospora beticola*)

30

Young sugar beet seedlings of the cultivar "ACH-31" were grown in pots to the 2 to 4 leaf stage. These plants were sprayed to run-  
off with an aqueous suspension, containing the concentration of active ingredient mentioned below, prepared from a stock solution  
35 containing 5 % of the active ingredient, 94 % cyclohexanone and 1 % emulsifier (Tween 20). After the plants had dried (3-5h), they were inoculated with a spore suspension of *Cercospora beticola* in an aqueous solution of 0.5 % gelatine. Then the trial plants were immediately transferred to a humid chamber with 18 to 23°C and a  
40 relative humidity close to 100 % and kept there for 5 days. For a period of further 10 to 14 days a cultivation in a greenhouse followed at 21 to 23°C and a relative humidity about 95 %. Then the extent of fungal attack on the leaves was visually assessed as % diseased leaf area.

45


In this test, the plants which had been treated with 200 ppm of compounds I-1 to I-4, I-6, I-8, and I-11 to I-19 showed an infection of up to 15%, whereas the untreated plants were infected to 85%.

5

### Comparison Trials

Active compounds A to E known from WO-A 98/46607, WO-A 98/46608, and WO-A 99/48893 resp., were used as comparison compounds in the following comparison trials:

15



20

No.	known from	R <sup>1</sup>	L <sub>n</sub>
A	WO-A 98/46607, Ex. 12	H	2,4,6-F <sub>3</sub>
B	WO-A 98/46608, Ex. 1	H	2-Cl-6-F
C	WO-A 98/46608, Ex. 5	CH <sub>3</sub>	2,6-F <sub>2</sub>
D	WO-A 99/48893, Ex. 6	H	2,6-F <sub>2</sub> -4-OCH <sub>3</sub>
E	WO-A 99/48893, Ex. 7	CH <sub>3</sub>	2,6-F <sub>2</sub> -4-OCH <sub>3</sub>

25

Comparison trial 1: Fungicidal control of rice blast caused by *Pyricularia oryzae* (protective)

30

Leaves of pot-grown rice seedling of the variety "Tai-Nong 67" were sprayed to run-off with an aqueous suspension, containing the concentration of active ingredient as described below prepared from a stock solution containing 10 % of the active ingredient, 85 % cyclohexanone and 5 % emulsifier. The plants were allowed to air-dry. At the following day the plants were inoculated with an aqueous spore suspension of *Pyricularia oryzae* containing  $1.0 \times 10^6$  spores/ml. Then the trial plants were immediately transferred to a humid chamber. After 6 days at 22-24°C and a relative humidity close to 100 % the extent of fungal attack on the leaves was visually assessed as % diseased leaf area.

40

In this trial, the plants which have been treated with 200 ppm of compounds I-1, I-2 and I-3, resp., showed an infection of 3 %, whereas the plants treated with 200 ppm of comparison compounds A, B and C, resp., were infected to 60, 10, and 20 %, resp., and the untreated plants were infected to 90 %.

45

Comparison trial 2: Control of powdery mildew on wheat caused by *Blumeria graminis f. sp. tritici*

The first fully developed leaves of pot grown wheat of the variety "Kanzler" were sprayed to run-off with an aqueous suspension, containing the concentration of active ingredient or their mixture mentioned in the table below, prepared from a stock solution containing 10 % of the active ingredient, 85 % cyclohexanone and 5 % emulsifier. The next day the treated plants were inoculated with spores of *Blumeria graminis f. sp. tritici* (= syn. *Erysiphe graminis f. sp. tritici*) by shaking heavily infested stock plants over the treated pots. After cultivation in the greenhouse for 7 days at 22-26°C and a relative humidity between 60 to 90 % the extent of fungal attack on the leaves was visually assessed as % diseased leaf area.

In this trial, the plants which have been treated with 200, and 50 ppm, resp. of compound I-7 showed an infection of 3, and 7 %, whereas the plants treated with 200, and 50 ppm, resp. of comparison compounds D and E, resp., were infected (at 200 ppm) to 60 and 80 %, resp., and (at 50 ppm) to 80 %, and the untreated plants were infected to 100 %.

25

30


35

40

45

## Claims:

1. 5-Halogen-6-phenyl-7-fluoroalkylamino-triazolopyrimidines of  
5 formula I



I

10 in which

15  $R^1$  is hydrogen, fluoro,  $C_1$ - $C_{10}$ -alkyl,  $C_2$ - $C_{10}$ -alkenyl,  $C_2$ - $C_{10}$ -alkynyl,  $C_2$ - $C_{10}$ -alkadienyl,

where the carbon chains of these radicals may be unsubstituted or partially or fully halogenated or may carry one to three groups  $R^a$ ,

20  $R^a$  is cyano, nitro, hydroxyl,  $C_1$ - $C_6$ -alkyl,  $C_1$ - $C_6$ -haloalkyl,  $C_1$ - $C_6$ -alkylcarbonyl,  $C_3$ - $C_6$ -cycloalkyl,  $C_1$ - $C_6$ -alkoxy,  $C_1$ - $C_6$ -haloalkoxy,  $C_1$ - $C_6$ -alkoxycarbonyl,  $C_1$ - $C_6$ -alkylthio,  $C_1$ - $C_6$ -alkylamino,

25 di- $C_1$ - $C_6$ -alkylamino,  $C_2$ - $C_6$ -alkenyl,  $C_2$ - $C_6$ -alkenyloxy,  $C_3$ - $C_6$ -alkynyloxy and  $C_1$ - $C_4$ -alkylenedioxy, which may be halogenated;

30  $R^2$  is hydrogen,  $C_1$ - $C_{10}$ -alkyl,  $C_2$ - $C_{10}$ -alkenyl,  $C_2$ - $C_{10}$ -alkynyl,  $C_2$ - $C_{10}$ -alkadienyl, where these radicals may be unsubstituted or partially or fully halogenated or may carry one to three groups  $R^a$ ,

35  $R^3$  is  $C_2$ - $C_8$ -fluoroalkyl or  $C_2$ - $C_8$ -fluoroalkenyl;


$X$  is halogen;

$n$  is 0 or an integer from 1 to 4;

40  $L$  each independently is halogen, nitro,  $C_1$ - $C_{10}$ -alkyl,  $C_1$ - $C_6$ -haloalkyl,  $C_1$ - $C_{10}$ -alkoxy or  $C_1$ - $C_6$ -haloalkoxy.

2. Compounds according to Claim 1 in which at least one  $L$  group  
45 is halogen.

3. Compounds according to Claims 1 or 2 in which R<sup>1</sup> is hydrogen, fluorine or methyl.
4. Compounds according to Claim 1 in which R<sup>2</sup> is hydrogen or C<sub>1</sub>-C<sub>10</sub>-alkyl.
5. Compounds according to claim 1 in which R<sup>3</sup> is a group A



wherein


15 Y<sup>1</sup> is hydrogen, fluoro or C<sub>1</sub>-C<sub>6</sub>-fluoroalkyl,

Y<sup>2</sup> is hydrogen or fluoro, or


Y<sup>1</sup> and Y<sup>2</sup> taken together form a double bond; and

20 m is 0 or 1.

6. A process for the preparation of compounds of formula I as defined in claim 1 which comprises reacting 5,7-dihalo-triazolopyrimidines of formula II



30 in which the variables have the meaning given in formula I with an amine of formula III



in which R<sup>1</sup>, R<sup>2</sup> and R<sup>3</sup> are defined as in formula I to give compounds of formula I.

7. A composition suitable for controlling phytopathogenic fungi, comprising a solid or liquid carrier and a compound of the formula I as claimed in claim 1.
8. A method for controlling phytopathogenic fungi, which comprises treating the fungi or the materials, plants, the soil or the seed to be protected against fungal attack with an effective amount of a compound of the formula I as claimed in claim 1.



# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 02/03829

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC 7 C07D487/04 A01N43/90		
According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) IPC 7 C07D		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) CHEM ABS Data, EPO-Internal, WPI Data, BEILSTEIN Data		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 46608 A (AMERICAN CYANAMID CO., USA) 22 October 1998 (1998-10-22) cited in the application claims 1,10	1-8
X	WO 98 46607 A (AMERICAN CYANAMID CO., USA) 22 October 1998 (1998-10-22) cited in the application claims 1,11; examples 34,40	1-8
<input type="checkbox"/> Further documents are listed in the continuation of box C. <span style="margin-left: 100px;"><input checked="" type="checkbox"/> Patent family members are listed in annex.</span>		
* Special categories of cited documents :		
<div style="display: flex; justify-content: space-between;"> <div style="width: 45%;"> <p>*A* document defining the general state of the art which is not considered to be of particular relevance</p> <p>*E* earlier document but published on or after the international filing date</p> <p>*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)</p> <p>*O* document referring to an oral disclosure, use, exhibition or other means</p> <p>*P* document published prior to the international filing date but later than the priority date claimed</p> </div> <div style="width: 45%;"> <p>*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention</p> <p>*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone</p> <p>*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.</p> <p>*G* document member of the same patent family</p> </div> </div>		
Date of the actual completion of the international search  <div style="text-align: center; font-weight: bold;">12 August 2002</div>		Date of mailing of the international search report  <div style="text-align: center; font-weight: bold;">21/08/2002</div>
Name and mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer  <div style="text-align: center; font-weight: bold;">Seelmann, I</div>

Form PCT/ISA/210 (second sheet) (July 1992)

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

Full/EP 02/03829

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9846608	A	22-10-1998	AU 735730 B2 12-07-2001
			AU 6867198 A 11-11-1998
			BG 103805 A 30-06-2000
			BR 9808531 A 23-05-2000
			CN 1257502 T 21-06-2000
			EE 9900486 A 15-06-2000
			EP 0975635 A1 02-02-2000
			HU 0001993 A2 28-10-2000
			JP 2001520650 T 30-10-2001
			NO 994973 A 13-10-1999
			NZ 500143 A 29-06-2001
			PL 336164 A1 05-06-2000
			SK 141499 A3 11-09-2001
			TR 9902552 T2 22-05-2000
			TW 460476 B 21-10-2001
			WO 9846608 A1 22-10-1998
			ZA 9803054 A 11-10-1999
WO 9846607	A	22-10-1998	AT 202779 T 15-07-2001
			AU 6576898 A 11-11-1998
			DE 69801048 D1 09-08-2001
			DE 69801048 T2 14-03-2002
			DK 975634 T3 24-09-2001
			EP 0975634 A1 02-02-2000
			ES 2160408 T3 01-11-2001
			PT 975634 T 28-12-2001
			WO 9846607 A1 22-10-1998
			ZA 9803055 A 11-10-1999

Form PCT/SA/210 (patent family annex) (July 1992)